



Clinical efficacy and safety of achieving very low LDL-cholesterol concentrations with the PCSK9 inhibitor evolocumab: a prespecified secondary analysis of the FOURIER trial

Robert P Giugliano, Terje R Pedersen, Jeong-Gun Park, Gaetano M De Ferrari, Zbigniew A Gaciong, Richard Ceska, Kalman Toth, Ioanna Gouni-Berthold, Jose Lopez-Miranda, François Schiele, François Mach, Brian R Ott, Estella Kanevsky, Armando Lira Pineda, Ransi Somaratne, Scott M Wasserman, Anthony C Keech, Peter S Sever, Marc S Sabatine, on behalf of the FOURIER Investigators

Summary

Background LDL cholesterol is a well established risk factor for atherosclerotic cardiovascular disease. How much one should or safely can lower this risk factor remains debated. We aimed to explore the relationship between progressively lower LDL-cholesterol concentrations achieved at 4 weeks and clinical efficacy and safety in the FOURIER trial of evolocumab, a monoclonal antibody to proprotein convertase subtilisin-kexin type 9 (PCSK9).

Methods In this prespecified secondary analysis of 25 982 patients from the randomised FOURIER trial, the relationship between achieved LDL-cholesterol concentration at 4 weeks and subsequent cardiovascular outcomes (primary endpoint was the composite of cardiovascular death, myocardial infarction, stroke, coronary revascularisation, or unstable angina; key secondary endpoint was the composite of cardiovascular death, myocardial infarction, or stroke) and ten prespecified safety events of interest was examined over a median of 2·2 years of follow-up. We used multivariable modelling to adjust for baseline factors associated with achieved LDL cholesterol. This trial is registered with ClinicalTrials.gov, number NCT01764633.

Findings Between Feb 8, 2013, and June 5, 2015, 27 564 patients were randomly assigned a treatment in the FOURIER study. 1025 (4%) patients did not have an LDL cholesterol measured at 4 weeks and 557 (2%) had already had a primary endpoint event or one of the ten prespecified safety events before the week-4 visit. From the remaining 25 982 patients (94% of those randomly assigned) 13 013 were assigned evolocumab and 12 969 were assigned placebo. 2669 (10%) of 25 982 patients achieved LDL-cholesterol concentrations of less than 0·5 mmol/L, 8003 (31%) patients achieved concentrations between 0·5 and less than 1·3 mmol/L, 3444 (13%) patients achieved concentrations between 1·3 and less than 1·8 mmol/L, 7471 (29%) patients achieved concentrations between 1·8 to less than 2·6 mmol/L, and 4395 (17%) patients achieved concentrations of 2·6 mmol/L or higher. There was a highly significant monotonic relationship between low LDL-cholesterol concentrations and lower risk of the primary and secondary efficacy composite endpoints extending to the bottom first percentile (LDL-cholesterol concentrations of less than 0·2 mmol/L; $p=0\cdot0012$ for the primary endpoint, $p=0\cdot0001$ for the secondary endpoint). Conversely, no significant association was observed between achieved LDL cholesterol and safety outcomes, either for all serious adverse events or any of the other nine prespecified safety events.

Interpretation There was a monotonic relationship between achieved LDL cholesterol and major cardiovascular outcomes down to LDL-cholesterol concentrations of less than 0·2 mmol/L. Conversely, there were no safety concerns with very low LDL-cholesterol concentrations over a median of 2·2 years. These data support further LDL-cholesterol lowering in patients with cardiovascular disease to well below current recommendations.

Funding Amgen.

Introduction

LDL cholesterol has been well established as a modifiable risk factor for atherosclerotic cardiovascular disease in epidemiological studies. In a series of landmark randomised controlled trials with statins, significant reductions in cardiovascular events were shown in patients with very high LDL-cholesterol concentrations (eg, decreasing LDL cholesterol from 5 to 3 mmol/L), average LDL-cholesterol concentrations (eg, decreasing LDL cholesterol from 3·5 to 2·5 mmol/L), and below

average LDL-cholesterol concentrations (eg, decreasing LDL cholesterol from 2·5 to 1·5–2 mmol/L).¹ Although the trials did not allocate patients to different LDL-cholesterol targets, the data suggested that lowering LDL cholesterol across a broad range of concentrations conferred similar cardiovascular risk reduction per unit reduction of LDL cholesterol. Based on these trials, the LDL-cholesterol target or threshold for treatment in published guidelines decreased to 2·5 mmol/L and then 1·8 mmol/L in high-risk patients.^{2–5}

Published Online
August 28, 2017
[http://dx.doi.org/10.1016/S0140-6736\(17\)32290-0](http://dx.doi.org/10.1016/S0140-6736(17)32290-0)

See Online/Comment
[http://dx.doi.org/10.1016/S0140-6736\(17\)32292-4](http://dx.doi.org/10.1016/S0140-6736(17)32292-4)

TIMI Study Group, Division of Cardiovascular Medicine, Brigham and Women's Hospital, Boston, MA, USA (R P Giugliano MD, J-Gun Park PhD, E Kanevsky MS, Prof M S Sabatine MD); Oslo University Hospital, Ullevål and Medical Faculty, University of Oslo, Oslo, Norway (Prof T R Pedersen MD); Department of Molecular Medicine, University of Pavia and Cardiac Intensive Care Unit and Laboratories for Experimental Cardiology, IRCCS Fondazione Policlinico San Matteo, Pavia, Italy (G M De Ferrari MD); Department of Internal Medicine, Hypertension and Vascular Diseases, The Medical University of Warsaw, Warsaw, Poland (Prof Z A Gaciong MD); Center of Preventive Cardiology, 3rd Department Internal Medicine, University General Hospital and 1st Medical Faculty, Prague, Czech Republic (Prof R Ceska MD); 1st Department of Medicine, University of Pécs, Pécs, Hungary (Prof K Toth MD); Polyclinic for Endocrinology, Diabetes, and Preventive Medicine, University of Cologne, Cologne, Germany (Prof I Gouni-Berthold MD); Lipids and Atherosclerosis Unit, Maimonides Biomedical Research Institute of Cordoba, Reina Sofia University Hospital, University of Cordoba, CIBEROBN, Cordoba, Spain (Prof J Lopez-Miranda MD); University Hospital Center Besançon, Besançon, France

(Prof F Schiele MD); Hôpital Cantonal, Hôpitaux Universitaires de Genève, Geneva, Switzerland

(Prof F Mach MD); Rhode Island Hospital, Department of Neurology, Alpert Medical School of Brown University, Providence, RI, USA

(Prof B R Ott MD); Amgen, Thousand Oaks, CA, USA

(A Lira Pineda MD, R Somaratne MD,

S M Wasserman MD); Sydney Medical School, National Health and Medical Research Council Clinical Trials Centre, University of Sydney, Sydney, NSW, Australia

(Prof A C Keech FRACP); and International Centre for Circulatory Health, National Heart and Lung Institute, Imperial College London, London, UK

(Prof P S Sever FRCP)

Correspondence to:

Dr Robert P Giugliano, Cardiovascular Division, Brigham and Women's Hospital, Boston, MA 02115, USA
rgiugliano@bwh.harvard.edu

Research in context

Evidence before this study

We searched MEDLINE on July 24, 2017, with the terms “cholesterol, LDL” and either “myocardial infarction” or “stroke.” The search was limited to publications from 2014 onwards since that was the date of a comprehensive meta-analysis on this topic. Abstracts were reviewed by two of the authors (RPG and MSS) to find publications describing the association of on-treatment LDL cholesterol and cardiovascular outcomes in patients with atherosclerotic cardiovascular disease. Relevant publications were supplemented with additional relevant publications known by the authors. None of the trials studying patients treated with statins provided data on a cutpoint of less than 1.3 mmol/L. We have published data on ezetimibe from IMPROVE-IT, in which the lowest cutpoint was 0.8 mmol/L, but there were fewer than 1000 patients in that subgroup. There is a published pooled analysis of smaller lipid-lowering trials of another proprotein convertase subtilisin-kexin type 9 inhibitor, alirocumab, but with approximately a tenth of the number of patients with an LDL-cholesterol concentration of less than 0.5 mmol/L.

Added value of this study

We found a strong relationship between achieved LDL cholesterol down to concentrations 0.2 mmol/L and a progressive reduction in major cardiovascular outcomes, with no increase in safety events. These observations extend previous findings with statins and ezetimibe to lower concentrations of LDL cholesterol than previously reported, in a larger sample size, and with the newest and most potent lipid-lowering therapy approved to date.

Implications of all the available evidence

All evidence to date from trials of intensive lipid lowering supports reduction of LDL cholesterol in high-risk patients to concentrations below those currently recommended in cholesterol guidelines. Studies with a longer follow-up period than in this study are needed to exclude the development of late complications of persistent very-low concentrations of LDL cholesterol.

Findings from the Further cardiovascular Outcomes Research with PCSK9 Inhibition in subjects with Elevated Risk (FOURIER) trial⁶ showed that the proprotein convertase subtilisin-kexin type 9 (PCSK9) monoclonal antibody evolocumab, when added to background statin therapy, lowered LDL-cholesterol concentrations to a median of 0.8 mmol/L (IQR 0.5–1.2) and significantly reduced the risk of cardiovascular events in patients with stable atherosclerotic cardiovascular disease who were followed up for a median of 2.2 years. No significant differences were found in major safety events or in prospective cognitive function testing between treatment groups.⁷

Notably, the LDL-cholesterol concentrations achieved were substantially lower than those in previous clinical outcome trials with lipid-lowering therapies, and it is not known whether there is a threshold below which there is no added clinical benefit, or whether there is an increase in adverse safety events. Our aim was to explore the relationship between progressively lower LDL cholesterol achieved at 4 weeks and clinical efficacy and safety in this prespecified, secondary analysis of the FOURIER trial.

Methods

Study design and participants

FOURIER⁸ was a randomised, double-blind, placebo-controlled trial that enrolled 27564 patients aged 40–85 years with stable atherosclerotic cardiovascular disease (previous myocardial infarction, previous non-haemorrhagic stroke, or symptomatic peripheral arterial disease) and additional risk factors placing them at increased cardiovascular risk. Eligible patients had LDL-cholesterol concentrations of 1.8 mmol/L or higher or non-HDL concentrations of 2.6 mmol/L or higher while

taking an optimised lipid-lowering regimen including a high or moderate intensity statin, with or without ezetimibe.

Key exclusion criteria were recent myocardial infarction or stroke within 4 weeks, previous haemorrhagic stroke, estimated glomerular filtration rate of less than 20 mL/min per 1.73 m², New York Heart Association class III or IV heart failure or left ventricular ejection fraction of less than 30%, malignancy in the previous 10 years, and elevation of creatine kinase more than five times above normal or hepatic aminotransferase more than three times above normal. All patients provided written informed consent. The protocol was approved by ethics committees at each centre.

Procedures

Patients were randomly assigned (1:1) either subcutaneous evolocumab (either 140 mg every 2 weeks or 420 mg once per month, per patient preference) or matching placebo injections, and were followed up for a median of 2.2 years (IQR 1.8–2.5 years).

After randomisation, follow-up visits occurred at weeks 2, 4, 12, and every 12 weeks thereafter. Blood specimens were obtained and sent to a central core laboratory for analysis at weeks 4, 12, 24, and every 24 weeks thereafter. Lipid concentrations after randomisation were not made available to patients, investigators, study personnel, or the endpoint adjudicators. LDL cholesterol was calculated based on the Friedewald equation,¹¹ unless the calculated value was less than 1.03 mmol/L (40 mg/dL) or the measured triglycerides were higher than 4.52 mmol/L (400 mg/dL), in which case a direct measurement using ultracentrifugation was done. Patients with an

LDL-cholesterol assessment at week 4 who did not have a primary efficacy or prespecified safety event prior to the week-4 visit were included in this analysis.

Outcomes

The primary endpoint of the FOURIER trial was the composite of cardiovascular death, myocardial infarction, stroke, coronary revascularisation, or hospital admission for unstable angina. The key secondary endpoint was the composite of cardiovascular death, myocardial infarction, or stroke. An independent clinical endpoint committee adjudicated all efficacy endpoints as well as new-onset diabetes mellitus, without knowledge to randomisation assignment or post-randomisation lipid values.

There were ten prespecified safety endpoints: serious adverse events, adverse events leading to study drug discontinuation (excluding injection-site reactions because these were more frequent with evolocumab, as was the achievement of very low LDL cholesterol, which would confound the results), elevation in hepatic transaminase concentration of more than three times above normal, new or recurrent cancer (excluding non-melanoma skin cancer), cataract-related adverse events, elevation in creatine kinase concentration of more than five times above normal, haemorrhagic stroke, neurocognitive adverse events, new-onset diabetes mellitus, and non-cardiovascular death. Investigators reported adverse events as verbatim terms that were then mapped to the Medical Dictionary for Regulatory Activities (version 19.1) preferred terms with over reading by trained coders.

In patients who participated in the Evaluating PCSK9 Binding Antibody Influence on Cognitive Health in High Cardiovascular Risk Subjects (EBBINGHAUS) study,^{7,9} an embedded study within the FOURIER trial, we evaluated the primary endpoint of spatial working memory strategy index (executive function), the three secondary endpoints (spatial working memory between-errors score, working memory; paired associates learning score adjusted, episodic memory; and five-choice reaction time, psychomotor speed), and a global composite score (average of the four scores), as assessed using the Cambridge Neuropsychological Test Automated Battery (CANTAB) tool, across the five groups of achieved LDL cholesterol. CANTAB scores were analysed in 1154 patients who had baseline cognitive testing completed before or on the first day of study drug, and who did not have an efficacy or prespecified safety event prior to the measurement of the LDL cholesterol at 4 weeks.

Patients were also asked for their self-assessment of changes in memory and executive function (including the subdomains of planning, organisation, and divided attention) at the end of the trial compared with the start using a shortened version (23 questions) of the Everyday Cognition (ECog) self-assessment scales¹⁰ (appendix pp 2–3). Lower scores represented better function (range 1–5).

Statistical analysis

Patients were categorised into five prespecified subgroups based on the LDL cholesterol at 4 weeks, irrespective of treatment assignment: <0.5 mmol/L, 0.5 to less than 1.3 mmol/L, 1.3 to less than 1.8 mmol/L, 1.8 to less than 2.6 mmol/L, and 2.6 mmol/L or higher. We tested trends in baseline patient characteristics across achieved LDL-cholesterol groups using the Jonkheere-Terpstra trend test for continuous variables and Cochran-Armitage trend test for categorical variables. Independent predictors of achieving an LDL-cholesterol concentration of less than 0.5 mmol/L at 4 weeks (in addition to treatment assignment) were identified and served as covariates to generate multivariable adjusted hazard ratios (HRs) using Cox proportional hazard models^{12,13} or odds ratios using logistic regression models¹⁴ for outcomes of interest. These variables included the LDL-cholesterol concentration at baseline, age, sex, race, body-mass index, geographical region, and use of a P2Y12 inhibitor. We calculated adjusted p_{trend} values using Cox proportional hazard regression by testing the coefficient of the achieved LDL groups or the Cochran-Armitage trend test of proportions in achieved LDL groups, as appropriate. We compared Kaplan-Meier estimates with the log-rank test. We plotted the relationship between composite efficacy endpoints and achieved LDL cholesterol using a smoothing function applied to the averages of estimated event rates at each LDL level based on the adjusted Cox models. In a post-hoc analysis, we evaluated efficacy and safety outcomes in patients with LDL cholesterol of less than 0.4 mmol/L (15 mg/dL) and less than 0.26 mmol/L (10 mg/dL) compared with patients with LDL-cholesterol concentrations of 2.6 mmol/L (100 mg/dL) or higher.

The raw database was provided to the Thrombolysis in Myocardial Infarction (TIMI) study group, which did data analyses independently of the study sponsor. A two-sided p value of less than 0.05 was considered evidence of a significant effect; no adjustments were made for assessment of multiple endpoints.¹⁵ Analyses were done in SAS (version 9.4). The FOURIER (NCT01764633) and EBBINGHAUS (NCT02207634) trials are registered with ClinicalTrials.gov.

Role of the funding source

The funder collected and interpreted the data, and helped edit the manuscript. The executive committee (MSS, TRP, RPG, ACK, and PSS) had full access to all the data in the study and had final responsibility for the decision to submit for publication.

Results

Of 27564 patients who were randomly assigned either evolocumab or placebo in the FOURIER trial between Feb 8, 2013, and June 5, 2015, 1025 (4%) patients did not have an LDL cholesterol measured at 4 weeks and 557 (2%) had already had a primary endpoint event or one

For the Cambridge Neuropsychological Test Automated Battery tool see www.cambridgecognition.com

See Online for appendix




Figure 1: Distribution of achieved LDL-cholesterol concentrations at 4 weeks in patients who did not have a primary efficacy or prespecified safety event before the study

Red bars are evolocumab (median 0.8 mmol/L, IQR 0.5–1.2). Blue bars are placebo (median 2.2 mmol/L, IQR 1.9–2.7).




Figure 2: Median LDL-cholesterol concentrations categorised by achieved LDL-cholesterol concentration at 4 weeks

Time-weighted median LDL-cholesterol concentrations from week 4 onwards for the five groups categorised by LDL cholesterol at 4 weeks from lowest to highest were 0.5 mmol/L, 0.8 mmol/L, 1.7 mmol/L, 2.3 mmol/L, and 2.9 mmol/L.

	LDL-cholesterol concentration at 4 weeks					P _{trend}
	<0.5 mmol/L (n=2669)	0.5 to <1.3 mmol/L (n=8003)	1.3 to <1.8 mmol/L (n=3444)	1.8 to <2.6 mmol/L (n=7471)	≥2.6 mmol/L (n=4395)	
Demographics						
Age, years	63.6 (8.9)	62.7 (8.9)	62.0 (9.1)	62.7 (8.8)	61.4 (9.0)	<0.0001
Weight, kg	81.0 (15.0)	85.5 (17.3)	86.9 (18.6)	85.7 (17.5)	85.7 (17.3)	<0.0001
Body-mass index, kg/m ²	28.1 (4.1)	29.4 (5.0)	30.1 (5.7)	29.5 (5.2)	29.5 (5.1)	<0.0001
Men	2231 (84%)	6149 (77%)	2499 (73%)	5651 (76%)	3146 (72%)	<0.0001
White	2127 (80%)	6847 (86%)	2902 (84%)	6355 (85%)	3856 (88%)	<0.0001
Medical history						
Myocardial infarction	2167 (81%)	6475 (81%)	2757 (80%)	6138 (82%)	3555 (81%)	0.09
Non-haemorrhagic stroke	534 (20%)	1541 (19%)	657 (19%)	1406 (19%)	859 (20%)	0.70
Peripheral arterial disease	320 (12%)	1081 (14%)	470 (14%)	903 (12%)	632 (14%)	0.0012
Hypertension	2081 (78%)	6393 (80%)	2808 (82%)	5974 (80%)	3561 (81%)	0.0051
Diabetes	944 (35%)	2912 (36%)	1478 (43%)	2637 (35%)	1488 (34%)	<0.0001
Current smoker	692 (26%)	2159 (27%)	999 (29%)	2093 (28%)	1410 (32%)	<0.0001
TIMI risk score for secondary prevention	3.18 (1.19)	3.25 (1.25)	3.38 (1.23)	3.26 (1.23)	3.35 (1.24)	<0.0001
Treatment at baseline						
High-intensity statin*	1685 (63%)	5548 (69%)	2406 (70%)	5225 (70%)	3149 (72%)	<0.0001
Ezetimibe	109 (4.1%)	403 (5.0%)	187 (5.4%)	347 (4.6%)	325 (7.4%)	<0.0001
Aspirin or P ₂ Y ₁₂ inhibitor	2496 (94%)	7398 (92%)	3175 (92%)	6917 (93%)	4020 (91%)	0.0290
Beta blocker	1993 (75%)	6088 (76%)	2620 (76%)	5640 (75%)	3303 (75%)	0.53
ACE-I or ARB, aldosterone antagonist, or both	2082 (78%)	6260 (78%)	2702 (78%)	5880 (79%)	3405 (77%)	0.61
Baseline laboratory measurements						
LDL cholesterol, mmol/L	2.1 (1.9–2.4)	2.4 (2.1–2.7)	2.2 (1.9–2.7)	2.3 (2.1–2.6)	3.0 (2.6–3.5)	<0.0001
Total cholesterol, mmol/L	4.0 (3.7–4.5)	4.3 (3.9–4.8)	4.2 (3.7–4.8)	4.2 (3.9–4.6)	5.0 (4.5–5.6)	<0.0001
HDL cholesterol, mmol/L	1.1 (0.9–1.3)	1.1 (1.0–1.3)	1.1 (0.9–1.3)	1.1 (1.0–1.4)	1.2 (1.0–1.4)	<0.0001
Triglycerides, mmol/L	1.5 (1.2–2.1)	1.5 (1.1–2.0)	1.6 (1.1–2.2)	1.4 (1.1–1.9)	1.6 (1.2, 2.1)	<0.0001
Lipoprotein(a), nmol/L	22 (9–53)	43 (14–176)	32 (12–153)	37 (13–163)	49 (16–188)	<0.0001

Data are mean (SD), n (%), or median (IQR). LDL-cholesterol values are rounded to the nearest 0.1 in mmol/L; the precise values converted from mg/dL are <0.52 mmol/L (<20 mg/dL), 0.52 to <1.29 mmol/L (20 to <50 mg/dL), 1.29 to <1.81 mmol/L (50 to <70 mg/dL), 1.81 to <2.59 mmol/L (70 to <100 mg/dL), and ≥2.59 mmol/L (≥100 mg/dL). TIMI=Thrombolysis in Myocardial Infarction study group. ACE-I=angiotensin-converting-enzyme inhibitor. ARB=angiotensin receptor blocker. *Defined as having an LDL-lowering potency equal to or greater than that of atorvastatin 40 mg daily.

Table 1: Baseline characteristics by achieved LDL-cholesterol concentration at 4 weeks after randomisation

of the ten prespecified safety events before the week-4 visit. There were no differences between treatment groups in the number of patients who did not have a week-4 LDL cholesterol measurement or in patients who had a clinical event before 4 weeks. The final analysis set consisted of 25 982 patients (94% of those included at randomisation) of whom 13 013 were randomly assigned evolocumab and 12 969 were randomly assigned placebo. 2669 (10%) patients enrolled in FOURIER achieved an LDL cholesterol of less than 0.5 mmol/L at 4 weeks (figure 1). Of these 2669 patients, the median LDL cholesterol at 4 weeks was 0.36 mmol/L (IQR 0.28–0.44). The median LDL cholesterol over time for each of the five groups categorised by the LDL-cholesterol concentration at week 4 is shown in figure 2. Across the five groups of achieved LDL cholesterol at 4 weeks ordered from lowest to highest concentration groups, 2659 (99.6%) of 2669 patients, 7721 (96%) of 8003 patients, 1427 (41%) of 3444 patients, 782 (10%) of 7471 patients, and

424 (10%) of 4395 patients were randomly assigned evolocumab. The rates of study drug persistence were high during the trial (about 90% at 2 years) and similar across the five groups categorised by achieved LDL-cholesterol concentration at 4 weeks (appendix p 4). Baseline characteristics varied across the achieved LDL-cholesterol categories at week 4 (table 1).

The risk of the primary composite efficacy endpoint after week 4 was lower with decreasing achieved LDL-cholesterol concentrations at week 4 (figure 3A). The Kaplan-Meier event rates at 3 years across the five groups of achieved LDL-cholesterol concentrations from lowest to highest groups were 10.3%, 12.4%, 13.6%, 13.7%, and 15.5% ($p_{\text{trend}} < 0.0001$). The corresponding adjusted HRs using the group with LDL cholesterol of 2.6 mmol/L or greater as the reference group were 0.76 (95% CI 0.64–0.90), 0.85 (0.76–0.96), 0.94 (0.82–1.09), and 0.97 (0.86–1.09), for the next four groups from the lowest to the highest achieved LDL cholesterol




Figure 3: Relationship between the achieved LDL-cholesterol concentration at 4 weeks and the risk of the primary (A) and key secondary (B) efficacy composite endpoints

The primary efficacy endpoint was a composite of cardiovascular death, myocardial infarction, stroke, coronary revascularisation, or hospital admission for unstable angina. The key secondary efficacy endpoint was a composite of cardiovascular death, myocardial infarction, or stroke. The red line represents the adjusted probability of an event and blue areas are the 95% CIs of the regression model estimate.

($p_{\text{trend}} < 0.0001$; appendix pp 5–6). Regression modelling of the relationship between achieved LDL cholesterol at 4 weeks and the subsequent risk of cardiovascular events showed a steady decline in risk as achieved LDL-cholesterol

concentration decreased ($p = 0.0012$ for the β coefficient), even down to the first percentile (LDL cholesterol < 0.2 mmol/L, figure 3A). A similar monotonic reduction in the key secondary endpoint was observed (figure 3B,

	LDL-cholesterol concentration at 4 weeks					P _{trend}
	<0.5 mmol/L (n=2669)	0.5 to <1.3 mmol/L (n=8003)	1.3 to <1.8 mmol/L (n=3444)	1.8 to <2.6 mmol/L (n=7471)	≥2.6 mmol/L (n=4395)	
Serious adverse events	614 (23%)	1948 (24%)	838 (24%)	1684 (23%)	1022 (23%)	0.13
Adjusted OR (95% CI)	0.97 (0.86–1.10)	1.01 (0.92–1.11)	1.01 (0.90–1.13)	0.93 (0.84–1.02)	1 (ref)	0.30
Adverse events* leading to discontinuation of study drug	98 (4%)	295 (4%)	124 (4%)	234 (3%)	149 (3%)	0.11
Adjusted OR (95% CI)	1.08 (0.82–1.43)	1.07 (0.86–1.33)	1.07 (0.83–1.39)	0.91 (0.73–1.14)	1 (ref)	0.13
AST or ALT elevation (>3 times ULN)	41 (2%)	120 (1%)	76 (2%)	119 (2%)	83 (2%)	0.19
Adjusted OR (95% CI)	0.96 (0.64–1.43)	0.87 (0.64–1.17)	1.25 (0.90–1.74)	0.91 (0.68–1.24)	1 (ref)	0.64
Creatine kinase elevation (>5 times ULN)	18 (1%)	55 (1%)	19 (1%)	58 (1%)	26 (1%)	0.99
Adjusted OR (95% CI)	1.02 (0.53–1.96)	1.07 (0.65–1.77)	0.88 (0.47–1.65)	1.23 (0.75–2.02)	1 (ref)	0.72
Neurocognitive events	49 (2%)	122 (2%)	51 (1%)	100 (1%)	52 (1%)	0.019
Adjusted OR (95% CI)	1.28 (0.84–1.96)	1.10 (0.78–1.55)	1.10 (0.73–1.65)	0.97 (0.68–1.39)	1 (ref)	0.15
New onset diabetes mellitus†	135/1655 (8%)	389/4863 (8%)	162/1886 (9%)	356/4603 (8%)	220/2778 (8%)	0.63
Adjusted OR (95% CI)	1.06 (0.83–1.35)	1.00 (0.83–1.20)	1.03 (0.83–1.30)	0.95 (0.78–1.14)	1 (ref)	0.48
Cataract-related adverse events	56 (2%)	124 (2%)	61 (2%)	134 (2%)	55 (1%)	0.15
Adjusted OR (95% CI)	1.54 (1.03–2.31)	1.14 (0.82–1.60)	1.34 (0.91–1.98)	1.35 (0.96–1.89)	1 (ref)	0.43
New or progressive malignancy	64 (2%)	205 (3%)	87 (3%)	166 (2%)	99 (2%)	0.22
Adjusted OR (95% CI)	0.90 (0.64–1.27)	1.01 (0.78–1.31)	1.04 (0.77–1.42)	0.88 (0.67–1.15)	1 (ref)	0.72
Haemorrhagic stroke	3 (<1%)	19 (<1%)	7 (<1%)	17 (<1%)	7 (<1%)	0.99
Adjusted HR (95% CI)	0.71 (0.17–2.90)	1.55 (0.62–3.85)	1.39 (0.47–4.14)	1.57 (0.62–3.98)	1 (ref)	0.91
Non-cardiovascular death	25 (1%)	86 (1%)	34 (1%)	66 (1%)	45 (1%)	0.67
Adjusted HR (95% CI)	0.89 (0.53–1.50)	1.06 (0.72–1.55)	1.03 (0.65–1.64)	0.89 (0.60–1.33)	1 (ref)	0.73

Data are n (%) or n/N (%), unless otherwise specified. OR=odds ratio. ref=reference. AST=aspartate aminotransferase. ALT=alanine aminotransferase. ULN=upper limit of normal. HR=hazard ratio.
*Excludes 17 patients with injection-site reactions. †Denominator excludes patients who were diagnosed with diabetes mellitus before the week-4 visit.

Table 2: Safety events by achieved LDL-cholesterol concentration at 4 weeks after randomisation

$p=0.0001$ for the β coefficient), with the group achieving an LDL-cholesterol concentration of less than 0.5 mmol/L at 4 weeks having the lowest risk of cardiovascular death, myocardial infarction, or stroke (3-year Kaplan-Meier rate 6.6%) with an adjusted HR of 0.69 (95% CI 0.56–0.85) compared with the group with LDL-cholesterol concentrations of 2.6 mmol/L or higher (appendix p 5). These findings were associated with reductions in myocardial infarction, ischaemic stroke, and coronary revascularisation (appendix pp 5–6, with no apparent relationship between achieved LDL cholesterol and unstable angina, cardiovascular death, or all-cause mortality in the timeframe studied).

Serious adverse events after week 4 occurred in 6106 (24%) patients. Less than 4% of all patients had an adverse event leading to drug discontinuation, and there were no differences in these events by achieved LDL-cholesterol concentration at 4 weeks (table 2). No differences in elevations in hepatic aminotransferases, creatine kinase, new-onset diabetes mellitus, cataract-related adverse events, new or progressive malignancy, haemorrhagic stroke, or non-cardiovascular death were observed. In an unadjusted analysis, more patients had neurocognitive events who had lower LDL cholesterol at 4 weeks ($p_{\text{trend}}=0.019$); however, after adjustment for differences in baseline characteristics, the association was no longer significant (adjusted $p_{\text{trend}}=0.15$; table 2).

To further explore the relationship between achieved LDL-cholesterol concentration and cognition, we also analysed 1154 patients who had formal cognitive testing as part of the EBBINGHAUS study, on or before the day of the first dose of study drug. No differences across the five groups of achieved LDL-cholesterol concentration at 4 weeks were observed in the change from baseline in the primary cognitive endpoint of spatial working memory strategy index of executive function, in any of the three secondary endpoints, or in the global composite score (appendix pp 7–8).

Patient-reported changes in memory and three sub-domains of executive function from the start to the end of the trial are shown in the appendix (p 9). The total score was significantly better in patients with lower achieved LDL cholesterol ($p_{\text{trend}}=0.0168$), although the absolute differences in mean scores were small and the relationship across LDL-cholesterol groups was not monotonic.

In a post-hoc analysis of patients with ultra-low LDL cholesterol, 1335 (5%) patients achieved an LDL-cholesterol concentration of less than 0.4 mmol/L (15 mg/dL) and 504 (2%) patients achieved an LDL-cholesterol concentration of less than 0.26 mmol/L (10 mg/dL) at week 4 (appendix p 10). The median LDL-cholesterol concentration in the latter group was 0.18 mmol/L (7 mg/dL) with an IQR of 0.13–0.23 (5–9 mg/dL). Major

cardiovascular events progressively declined with decreasing achieved LDL-cholesterol concentrations, with adjusted HRs in the group with LDL-cholesterol concentrations of less than 0.26 mmol/L at 4 weeks of 0.69 (95% CI 0.49–0.97) for the primary and 0.59 (0.37–0.92) for the key secondary endpoints compared with the reference group (LDL cholesterol \geq 2.6 mmol/L). Neither serious adverse events nor adverse events leading to drug discontinuation occurred in excess in these two groups with ultra-low LDL cholesterol compared with patients in the reference group.

Discussion

In an analysis of over 25 000 patients with atherosclerotic cardiovascular disease, we found a strong relationship between on-treatment LDL-cholesterol concentration and major cardiovascular outcomes. Conversely, we found no significant association between LDL-cholesterol concentrations and prespecified adverse events in patients followed up for a median of 2.2 years.

These data are supported by previous observations from several types of analyses. First, subgroup analyses from clinical trials have showed consistent benefit with LDL-cholesterol lowering therapy even in subgroups with the lowest baseline concentrations of LDL cholesterol. Specifically, in the Cholesterol Treatment Trialists Collaboration, the clinical benefit of LDL-cholesterol lowering with statins was consistent even in patients whose baseline LDL-cholesterol concentration was less than 2 mmol/L, in whom statin therapy lowered LDL-cholesterol concentrations to less than approximately 1.6 mmol/L.¹ Likewise, in the JUPITER trial¹⁶ of primary prevention, there was a consistent clinical benefit with rosuvastatin even in patients starting with an LDL-cholesterol concentration of less than 1.6 mmol/L, in whom rosuvastatin lowered LDL-cholesterol concentrations to less than approximately 0.8 mmol/L. In the IMPROVE-IT trial,¹⁷ there was consistent benefit even in patients in the lowest quartile of LDL cholesterol at randomisation (median 1.8 mmol/L) in whom the LDL cholesterol was reduced to 1.2 mmol/L. Second, in analyses of on-treatment LDL cholesterol from clinical trials of statins, a monotonic relationship was observed between LDL-cholesterol concentration and cardiovascular events.^{18,19} However, the lowest subgroup in those analyses only extended down to less than 1.3 mmol/L. Using data from IMPROVE-IT,²⁰ we extended that relationship down to less than 0.8 mmol/L, but that subgroup consisted of fewer than 1000 patients.

We now extend these data to unprecedented low concentrations of LDL cholesterol, with the lowest prespecified group (2669 patients) with an on-treatment LDL-cholesterol concentration of less than 0.5 mmol/L, a median LDL cholesterol of 0.36 mmol/L, and a quarter of that subgroup with an LDL-cholesterol concentration of less than 0.28 mmol/L, and an exploratory subgroup of 504 patients with an on-treatment LDL cholesterol of less

than 0.26 mmol/L (median 0.18 mmol/L). These clinical data are supported by mechanistic imaging data from the GLAGOV study²¹ showing a similar monotonic relationship between on-treatment LDL-cholesterol concentrations and coronary atheroma volume regression down to 0.5 mmol/L. Our observations are also supported by data from a pooled analysis of smaller lipid-lowering trials of another PCSK9 inhibitor, alirocumab.²² Establishment of which pharmacotherapies to use to achieve these concentrations of LDL cholesterol will require careful consideration in different health-care systems.

The association between LDL cholesterol and cardiovascular events was strongest for myocardial infarction, stroke, and coronary revascularisation, and not seen for cardiovascular death. It has been long-recognised, since the Cholesterol Treatment Trialists Collaboration, that the relative risk reduction per unit LDL-cholesterol lowering with statins is less for vascular death than for myocardial infarction or stroke.¹ It has also been observed in clinical trials that a mortality benefit from LDL-cholesterol lowering typically takes years to emerge,^{23,24} and the median duration of follow-up in FOURIER was only 2.2 years.

With regard to safety, we found similar adjusted risk of ten prespecified safety events regardless of the achieved LDL-cholesterol concentration, even in the 2669 patients who achieved an LDL-cholesterol concentration of less than 0.5 mmol/L at 4 weeks. These findings are unique in that they represent the first analysis of a large cohort of patients to achieve such very low LDL-cholesterol concentrations, namely being less than a third of the most common treatment goal (<1.8 mmol/L) for the highest-risk patients.^{25,26}

Our finding of no increase in adverse events in patients achieving very low LDL-cholesterol concentrations with the combination of evolocumab plus a statin is similar to that reported in the IMPROVE-IT trial²⁰ with ezetimibe plus simvastatin, in which no excess in adverse safety events were reported even in patients who achieved an LDL-cholesterol concentration of less than 0.8 mmol/L at 4 weeks and who were followed up for an average of 6 years. Both IMPROVE-IT and FOURIER were randomised trials of newer therapies on a background of a statin. In analyses from both of these trials, no significant increase in risk in adverse events such as those that had been reported in earlier placebo-controlled trials of statins (eg, new-onset diabetes mellitus,²⁶ elevations in hepatic transaminases,²⁷ and muscle safety²⁸) were observed. Additionally, neither the FOURIER study nor the EBBINGHAUS study reported increases in neurocognitive events, as reported in a meta-analysis²⁹ of phase 2 and 3 trials of PCSK9 inhibitors, or cataract-related events, as reported in 839 patients who achieved LDL-cholesterol concentrations of less than 0.6 mmol/L drawn from 14 randomised controlled trials with alirocumab.³⁰

The largest evaluation of the safety of evolocumab before FOURIER showed a favourable safety profile and

good tolerability up until 4 years,³¹ but this pooled analysis of four phase 2 trials was limited by a modest size (n=1324) and use of an open-label design comparing evolocumab with standard of care during the extension phase. In the FOURIER trial, because virtually all of the 2669 patients who achieved an LDL-cholesterol concentration of less than 0.5 mmol/L at 4 weeks had been randomly assigned evolocumab, the data from our current analysis now adds to the evidence supporting the safety of evolocumab, even in patients who achieve very low LDL-cholesterol concentrations.

The major limitations of this analysis were the absence of randomisation to on-treatment LDL-cholesterol concentrations (patients were classified based on post-randomisation measurement of LDL cholesterol), a median follow-up of 2.2 years, and low frequency of some safety events. We used multivariable adjustment to limit confounding due to differences in baseline characteristics across the groups of achieved LDL cholesterol, but acknowledge that this might have been incomplete and one cannot adjust for unmeasured confounders. Although most safety endpoints evaluated would be expected to manifest within the 2.2 years of follow-up based on experiences with other lipid therapies,^{27,28} we acknowledge that adverse events such as cancer can take longer to manifest. However, in a smaller open-label extension study³¹ of evolocumab with 4 years of follow-up, no safety concerns have emerged, and two long-term extension studies of FOURIER following approximately 6600 patients (NCT03080935 and NCT02867813) that are planned to last 5 years should provide longer-term insights. In conclusion, we observed that patients who achieved progressively lower LDL-cholesterol concentrations at 4 weeks in the FOURIER trial had progressively fewer cardiovascular events with no evidence of a plateau and with no increase in adverse events. These data support the use of intensive lipid-lowering therapies to prevent recurrent cardiovascular events in high-risk patients and suggest that a lower target LDL cholesterol than recommended in current guidelines (eg, <0.5 mmol/L) can safely be considered for the highest-risk patients.

Contributors

RPG did the literature search, designed the study, interpreted the data, wrote the first and subsequent drafts, and constructed the figures. TRP, GMDF, FM, BRO, ACK, and PSS designed the study, interpreted the data, and edited the manuscript. J-GP did data analysis, data interpretation, editing, and designed the figures. ZAG, RC, KT, IG-B, JL-M, and FS interpreted the data and edited the manuscript. EK did the data analysis, data interpretation, and editing. ALP and RS did data interpretation and editing. SMW designed the study, collected and interpreted the data, and edited the manuscript. MSS did a literature search, designed the study, interpreted the data, and edited and wrote part of the manuscript.

Declaration of interests

RPG reports grants and personal fees from Amgen, during the conduct of the study; grants and personal fees from Amgen, Daiichi-Sankyo, and Merck; and personal fees from Amgen, American College of Cardiology, Angel Med, Beckman-Coulter, Boehringer Ingelheim, Bristol-Myers Squibb, CVS Caremark, GlaxoSmithKline, Janssen, Lexicon, Portola, Pfizer, Regeneron, Sanofi-Aventis, St Jude, and Stealth Peptides, outside from the submitted work. TRP reports grants and personal fees from Amgen, during the conduct of the study; and personal fees from Amgen, Sanofi,

Merck, Boehringer-Ingelheim, and The Medicines Company, outside the submitted work. GMDF reports grants and personal fees from Amgen, during the conduct of the study; grants and personal fees from Amgen and MSD; grants from Boston Scientific; and personal fees from LivaNova and Sima Tau, outside the submitted work. ZAG reports personal fees from Amgen and Sanofi, during the conduct of the study. RC reports grants from Pfizer-International Atherosclerosis Society; and personal fees from MSD, Bayer, Amgen, Boehringer Ingelheim, Sanofi, AstraZeneca, NovoNordisk, Servier, and Kowa, outside the submitted work. IG-B reports personal fees and non-financial support from Amgen and Sanofi; and personal fees from Eli Lilly, Regeneron, and Aegerion, outside the submitted work. JL-M reports personal fees and non-financial support from Amgen and Sanofi; personal fees from MSD and Laboratorios Dr Esteve, outside the submitted work. FM reports grants, personal fees, and non-financial support from Amgen, MSD, Sanofi, AstraZeneca, and Pfizer, during the conduct of the study. BRO reports personal fees from Amgen, during the conduct of the study; participation on a data safety monitoring board for Accera; and grants from Long Term Care Group, Merck, Eli Lilly, TauRx, Janssen, Biogen, Avid, and Hoffman-La Roche, outside the submitted work. EK reports grants from Amgen, during the conduct of the study; and grants from Amgen and AstraZeneca, outside the submitted work. ALP is an employee of and holds stocks in Amgen. RS is an employee of Amgen and, as such, has received salary, bonus, stock or stock options, health insurance, and benefits; and is identified as an inventor on at least one pending patent application owned by Amgen relating to evolocumab. SMW is an employee of Amgen; and has a patent for evolocumab issued to Amgen. ACK reports grants and personal fees from Abbott, and Mylan; and personal fees from Amgen, AstraZeneca, and Pfizer, outside the submitted work. PSS reports grants and personal fees from Amgen, during the conduct of the study; grants and personal fees from Pfizer, outside the submitted work; and that he is the recipient of a National Institute for Health Research Senior Investigator Award and receives support from the Biomedical Research Centre Award to Imperial College Healthcare NHS Trust. MSS reports grants from Abbott Laboratories, Clinical Diagnostics, Daiichi-Sankyo, Gilead, GlaxoSmithKline, Roche Diagnostics, Takeda, Novartis, Poxel, Eisai, Genzyme, and Pfizer; grants and personal fees from Amgen, AstraZeneca, Intarcia, Merck, Janssen Research Development, The Medicines Company, and MedImmune; and personal fees from Alnylam, CVS Caremark, Lonis, Cubist, Esperion, and MyoKardia, outside the submitted work. J-GP, KT, and FS declare no competing interests.

References

- 1 Cholesterol Treatment Trialists Collaborators. Efficacy and safety of more intensive lowering of LDL cholesterol: a meta-analysis of data from 170 000 participants in 26 randomised trials. *Lancet* 2010; **376**: 1670–81.
- 2 Grundy SM, Cleeman JI, Merz CN, et al. Implications of recent clinical trials for the National Cholesterol Education Program Adult Treatment Panel III guidelines. *Circulation* 2004; **110**: 227–39.
- 3 Jacobson TA, Ito MK, Maki KC, et al. National Lipid Association recommendations for patient-centered management of dyslipidemia: part 1—full report. *J Clin Lipidol* 2015; **9**: 129–69.
- 4 Jacobson TA, Maki KC, Orringer CE, et al. National Lipid Association recommendations for patient-centered management of dyslipidemia: part 2. *J Clin Lipidol* 2015; **9** (suppl 6): S1–122.e121.
- 5 Catapano AL, Graham I, De Backer G, et al. 2016 ESC/EAS Guidelines for the management of dyslipidaemias. *Eur Heart J* 2016; **37**: 2999–3058.
- 6 Sabatine MS, Giugliano RP, Keech AC, et al. Evolocumab and clinical outcomes in patients with cardiovascular disease. *N Engl J Med* 2017; **376**: 1713–22.
- 7 Giugliano RP, Mach F, Zavitz K, et al. Evaluation of cognitive function in a randomized trial of evolocumab. *N Engl J Med* 2017; **377**: 633–43.
- 8 Sabatine MS, Giugliano RP, Keech A, et al. Rationale and design of the Further cardiovascular Outcomes Research with PCSK9 Inhibition in subjects with Elevated Risk trial. *Am Heart J* 2016; **173**: 94–101.
- 9 Giugliano RP, Mach F, Zavitz K, et al. Design and rationale of the EBBINGHAUS trial: a phase 3, double-blind, placebo-controlled, multicenter study to assess the effect of evolocumab on cognitive function in patients with clinically evident cardiovascular disease and receiving statin background lipid-lowering therapy—a cognitive study of patients enrolled in the FOURIER trial. *Clin Cardiol* 2017; **40**: 59–65.

- 10 Farias ST, Mungas D, Reed BR, et al. The measurement of everyday cognition (ECog): scale development and psychometric properties. *Neuropsychology* 2008; **22**: 531–44.
- 11 Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem* 1972; **18**: 499–502.
- 12 Concato J, Peduzzi P, Holford TR, Feinstein AR. Importance of events per independent variable in proportional hazards analysis. I. Background, goals, and general strategy. *J Clin Epidemiol* 1995; **48**: 1495–501.
- 13 Peduzzi P, Concato J, Feinstein AR, Holford TR. Importance of events per independent variable in proportional hazards regression analysis. II. Accuracy and precision of regression estimates. *J Clin Epidemiol* 1995; **48**: 1503–10.
- 14 Peduzzi P, Concato J, Kemper E, Holford TR, Feinstein AR. A simulation study of the number of events per variable in logistic regression analysis. *J Clin Epidemiol* 1996; **49**: 1373–79.
- 15 Rothman KJ. No adjustments are needed for multiple comparisons. *Epidemiology* 1990; **1**: 43–46.
- 16 Cannon CP, Blazing MA, Giugliano RP, et al. Ezetimibe added to statin therapy after acute coronary syndromes. *N Engl J Med* 2015; **372**: 2387–97.
- 17 Giugliano RP, Cannon CP, Blazing MA, et al. Baseline LDL-C and clinical outcomes with addition of ezetimibe to statin in 18,144 patients post ACS. *J Am Coll Cardiol* 2015; **65** (suppl 10): A4.
- 18 Boekholdt SM, Hovingh GK, Mora S, et al. Very low levels of atherogenic lipoproteins and the risk for cardiovascular events: a meta-analysis of statin trials. *J Am Coll Cardiol* 2014; **64**: 485–94.
- 19 Wiviott SD, Cannon CP, Morrow DA, et al. Can low-density lipoprotein be too low? The safety and efficacy of achieving very low low-density lipoprotein with intensive statin therapy: a PROVE IT-TIMI 22 substudy. *J Am Coll Cardiol* 2005; **46**: 1411–16.
- 20 Giugliano RP, Wiviott SD, Blazing MA, et al. Long-term safety and efficacy of achieving very low levels of low-density lipoprotein cholesterol: a prespecified analysis of the IMPROVE-IT trial. *JAMA Cardiol* 2017; **2**: 547–55.
- 21 Nicholls SJ, Puri R, Anderson T, et al. Effect of evolocumab on progression of coronary disease in statin-treated patients: the GLAGOV randomized clinical trial. *JAMA* 2016; **316**: 2373–84.
- 22 Ray K, Ginsberg H, Davidson M, et al. Relationship between percentage reduction in low-density lipoprotein cholesterol levels and major atherosclerotic cardiovascular disease among patients treated with statins +/- alirocumab or ezetimibe in the phase 3 ODYSSEY trials. *J Am Coll Cardiol* 2016; **67**: 2082.
- 23 Pedersen TR, Kjekshus J, Berg K, et al. Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S). *Lancet* 1994; **344**: 1383–89.
- 24 Long-Term Intervention with Pravastatin in Ischaemic Disease Study Group. Prevention of cardiovascular events and death with pravastatin in patients with coronary heart disease and a broad range of initial cholesterol levels. *N Engl J Med* 1998; **339**: 1349–57.
- 25 Orringer CE, Jacobson TA, Saseen JJ, et al. Update on the use of PCSK9 inhibitors in adults: recommendations from an expert panel of the National Lipid Association. *J Clin Lipidol* 2017; published online May 19. DOI:10.1016/j.jacl.2017.05.001.
- 26 Maki KC, Ridker PM, Brown WV, Grundy SM, Sattar N, The Diabetes Subpanel of the National Lipid Association Expert Panel. An assessment by the Statin Diabetes Safety Task Force: 2014 update. *J Clin Lipidol* 2014; **8** (suppl 3): S17–29.
- 27 Bays H, Cohen DE, Chalasani N, Harrison SA, The National Lipid Association's Statin Safety Task Force. An assessment by the Statin Liver Safety Task Force: 2014 update. *J Clin Lipidol* 2014; **8** (suppl 3): S47–57.
- 28 Rosenson RS, Baker SK, Jacobson TA, Kopecky SL, Parker BA, The National Lipid Association's Muscle Safety Expert Panel. An assessment by the Statin Muscle Safety Task Force: 2014 update. *J Clin Lipidol* 2014; **8** (suppl 3): S58–71.
- 29 Lipinski MJ, Benedetto U, Escarcega RO, et al. The impact of proprotein convertase subtilisin-kexin type 9 serine protease inhibitors on lipid levels and outcomes in patients with primary hypercholesterolaemia: a network meta-analysis. *Eur Heart J* 2016; **37**: 536–45.
- 30 Robinson JG, Rosenson RS, Farnier M, et al. Safety of very low low-density lipoprotein cholesterol levels with alirocumab: pooled data from randomized trials. *J Am Coll Cardiol* 2017; **69**: 471–82.
- 31 Koren MJ, Sabatine MS, Giugliano RP, et al. Long-term low-density lipoprotein cholesterol-lowering efficacy, persistence, and safety of evolocumab in treatment of hypercholesterolemia: results up to 4 years from the open-label OSLER-1 extension study. *JAMA Cardiol* 2017; **2**: 598–607.

THE LANCET

Supplementary appendix

This appendix formed part of the original submission and has been peer reviewed. We post it as supplied by the authors.

Supplement to: Giugliano RP, Pedersen TR, Park J-G, et al, on behalf of the FOURIER Investigators. Clinical efficacy and safety of achieving very low LDL-cholesterol concentrations with the PCSK9 inhibitor evolocumab (FOURIER trial): a prespecified secondary analysis. *Lancet* 2017; published online Aug 28. [http://dx.doi.org/10.1016/S0140-6736\(17\)32290-0](http://dx.doi.org/10.1016/S0140-6736(17)32290-0).

Supplemental Appendix – Table of Contents

Item	Title	Page #
Table S1.	Everyday Assessment of Cognition (Modified Short Version).....	2-3
Figure S2.	Percent of Patients Remaining on Study Drug Over Time Categorized by Achieved LDL-C at Four Weeks.....	4
Table S3.	Event Rates and Adjusted Risk of Efficacy Endpoints by Achieved LDL-C Group at Four Weeks.....	5-6
Table S4.	Changes From Baseline in the CANTAB Scores by Achieved LDL-C at Four Weeks.....	7-8
Table S5.	Patient Self-Report of Change in Memory and Executive Function Since the Start of the Trial.....	9
Table S6.	Primary and Key Secondary Endpoint and Adverse Events in Patients with Ultra-Low LDL-C at Four Weeks.....	10


Table S1. Everyday Cognition Questionnaire (Modified Short Version)

Directions: Please rate the subject's ability to perform certain everyday tasks **NOW**, as compared to his/her ability to do these same tasks at the beginning of the study. In other words, ask the study subject to try to remember how he/she was doing at the beginning of the study, and indicate any change he or she has seen. The amount of change is rated on a five-point scale ranging from: 1) no change or actually performs better than at the beginning of the study, 2) occasionally performs the task worse but not all of the time, 3) consistently performs the task a little worse than at the beginning of the study, 4) performs the task much worse than at the beginning of the study, or 5) don't know.

Compared to the beginning of the study, has there been any change in...	<i>Better or no change</i>	<i>Questionable / occasionally worse</i>	<i>Consistently a little worse</i>	<i>Consistently much worse</i>	<i>Don't know</i>
<i>Memory</i>					
1. Remembering a few shopping items without a list.	1	2	3	4	5
2. Remembering things that happened recently (such as recent outings, events in the news).	1	2	3	4	5
3. Recalling conversations a few days later.	1	2	3	4	5
4. Remembering where she/he has placed objects.	1	2	3	4	5
5. Repeating stories and/or questions.	1	2	3	4	5
6. Remembering the current date or day of the week.	1	2	3	4	5
7. Remembering he/she has already told someone something.	1	2	3	4	5
8. Remembering appointments, meetings, or engagements.	1	2	3	4	5
<i>Executive Functioning: Planning</i>					
1. Planning the sequence of stops on a shopping trip.	1	2	3	4	5
2. The ability to anticipate weather changes and plan accordingly (i.e. bring a coat or umbrella).	1	2	3	4	5
3. Developing a schedule in advance of anticipated events.	1	2	3	4	5
4. Thinking things through before acting.	1	2	3	4	5
5. Thinking ahead.	1	2	3	4	5

<i>Executive Functioning: Organization</i>					
1. Keeping living and work space organized.	1	2	3	4	5
2. Balancing the checkbook without error.	1	2	3	4	5
3. Keeping financial records organized.	1	2	3	4	5
4. Prioritizing tasks by importance.	1	2	3	4	5
5. Keeping mail and papers organized.	1	2	3	4	5
6. Using an organized strategy to manage a medication schedule involving multiple medications.	1	2	3	4	5
<i>Executive Functioning: Divided Attention</i>					
1. The ability to do two things at once.	1	2	3	4	5
2. Returning to a task after being interrupted.	1	2	3	4	5
3. The ability to concentrate on a task without being distracted by external things in the environment.	1	2	3	4	5
4. Cooking or working and talking at the same time.	1	2	3	4	5

Figure S2 - Percent of Patients Remaining on Study Drug Over Time Categorized by Achieved LDL-C at Four Weeks



LDL-C at 4 weeks	Number of Patients Still On Study Treatment							
<0.5 mmol/L	2662	2618	2570	2515	2461	1981	1440	880
0.5 to <1.3 mmol/L	7958	7792	7607	7455	7317	6064	4520	2772
1.3 to <1.8 mmol/L	3423	3337	3251	3166	3108	2493	1801	1039
1.8 to <2.6 mmol/L	7433	7285	7104	6955	6805	5605	4108	2495
≥ 2.6 mmol/L	4358	4262	4145	4043	3945	3260	2371	1491

Table S3 - Event Rates and Adjusted Risk of Efficacy Endpoints by Achieved LDL-C Group at Four Weeks

Endpoint	n (%)	Adjusted HR (95% CI)	P_{trend} of Adjusted HRs
Primary Efficacy Composite*			<0.0001
<0.5 mmol/L	217 (8.1)	0.76 (0.64-0.90)	
0.5-1.3 mmol/L	769 (9.6)	0.85 (0.76-0.96)	
1.3-1.8 mmol/L	354 (10.3)	0.94 (0.82-1.09)	
1.8-2.6 mmol/L	793 (10.6)	0.97 (0.86-1.09)	
≥ 2.6 mmol/L	521 (11.9)	referent	
CV death, MI, or stroke			<0.0001
<0.5 mmol/L	134 (5.0)	0.69 (0.56-0.85)	
0.5-1.3 mmol/L	455 (5.7)	0.75 (0.64-0.86)	
1.3-1.8 mmol/L	223 (6.5)	0.87 (0.73-1.04)	
1.8-2.6 mmol/L	498 (6.7)	0.90 (0.78-1.04)	
≥ 2.6 mmol/L	345 (7.8)	referent	
Cardiovascular death			0.83
<0.5 mmol/L	42 (1.6)	0.99 (0.67-1.47)	
0.5-1.3 mmol/L	140 (1.7)	1.07 (0.80-1.43)	
1.3-1.8 mmol/L	54 (1.6)	0.99 (0.69-1.43)	
1.8-2.6 mmol/L	134 (1.8)	1.14 (0.85-1.53)	
≥ 2.6 mmol/L	77 (1.8)	referent	
Myocardial infarction			<0.0001
<0.5 mmol/L	71 (2.7)	0.59 (0.45-0.78)	
0.5-1.3 mmol/L	266 (3.3)	0.69 (0.57-0.84)	
1.3-1.8 mmol/L	141 (4.1)	0.87 (0.69-1.09)	
1.8-2.6 mmol/L	297 (4.0)	0.85 (0.71-1.03)	
≥ 2.6 mmol/L	214 (4.9)	referent	
Stroke			0.0054
<0.5 mmol/L	45 (1.7)	0.81 (0.55-1.18)	
0.5-1.3 mmol/L	103 (1.3)	0.63 (0.47-0.85)	
1.3-1.8 mmol/L	56 (1.6)	0.81 (0.57-1.14)	
1.8-2.6 mmol/L	134 (1.8)	0.90 (0.68-1.20)	
≥ 2.6 mmol/L	92 (2.1)	referent	
Coronary revascularization			<0.0001
<0.5 mmol/L	111 (4.2)	0.63 (0.50-0.78)	
0.5-1.3 mmol/L	446 (5.6)	0.78 (0.67-0.91)	
1.3-1.8 mmol/L	215 (6.2)	0.91 (0.76-1.09)	
1.8-2.6 mmol/L	471 (6.3)	0.91 (0.78-1.05)	
≥ 2.6 mmol/L	326 (7.4)	referent	
Unstable angina			0.73
<0.5 mmol/L	44 (1.6)	1.18 (0.80-1.74)	
0.5-1.3 mmol/L	132 (1.6)	1.04 (0.78-1.39)	
1.3-1.8 mmol/L	51 (1.5)	0.95 (0.66-1.37)	
1.8-2.6 mmol/L	129 (1.7)	1.09 (0.82-1.47)	
≥ 2.6 mmol/L	80 (1.8)	referent	

All cause death				0.47
	<0.5 mmol/L	73 (2.7)	0.89 (0.66-1.20)	
	0.5-1.3 mmol/L	240 (3.0)	0.98 (0.79-1.23)	
	1.3-1.8 mmol/L	101 (2.9)	1.00 (0.77-1.31)	
	1.8-2.6 mmol/L	225 (3.0)	1.02 (0.82-1.27)	
	≥ 2.6 mmol/L	141 (3.2)	referent	

*Primary efficacy composite = cardiovascular death, MI, stroke, coronary revascularisation or hospitalization for unstable angina

CI = confidence interval, CV = cardiovascular, HR = hazard ratio

**Table S4. Changes From Baseline in CANTAB Scores by Achieved LDL-C at Four Weeks
(Patients in the Primary Analysis Set of the EBBINGHAUS Study)**

	Achieved LDL-C at Four Weeks (mmol/L)					P _{trend}
	<0.5 (N=107)	0.5-1.3 (N=356)	1.3-1.8 (N=156)	1.8-2.6 (N=326)	≥ 2.6 (N=209)	
Primary Endpoint: Spatial Working Memory Strategy Index of Executive Function						
Adjusted Z-score, mean (SE)	0.03 (0.08)	0.01 (0.05)	0.08 (0.06)	0.17 (0.05)	0.15 (0.06)	0.11
Difference from referent group, mean (95% CI)	-0.12 (-0.33, 0.09)	-0.14 (-0.31, 0.03)	-0.07 (-0.22, 0.08)	0.02 (-0.10, 0.14)	referent	
P-value compared to referent	0.25	0.11	0.34	0.73	referent	
Secondary Endpoints						
Spatial working memory between-errors score						
Adjusted Z-score, mean (SE)	0.03 (0.07)	0.04 (0.05)	0.08 (0.06)	0.09 (0.05)	0.08 (0.06)	0.61
Difference from referent group, mean (95% CI)	-0.05 (-0.25, 0.16)	-0.04 (-0.21, 0.12)	0.01 (-0.14, 0.15)	0.02 (-0.10, 0.13)	referent	
P-value compared to referent	0.66	0.62	0.94	0.80	referent	
Paired association learning score adjusted						
Adjusted Z-score, mean (SE)	0.10 (0.07)	0.09 (0.05)	0.11 (0.05)	0.16 (0.04)	0.12 (0.06)	0.61
Difference from referent group, mean (95% CI)	-0.02 (-0.20, 0.16)	-0.03 (-0.18, 0.12)	-0.01 (-0.14, 0.12)	0.04 (-0.06, 0.15)	referent	
P-value compared to referent	0.80	0.69	0.93	0.40	referent	
Reaction time 5-choice						
Adjusted Z-score, mean (SE)	-0.10 (0.08)	-0.10 (0.05)	0.01 (0.06)	0.02 (0.05)	-0.04 (0.06)	0.47
Difference from referent group, mean (95% CI)	-0.07 (-0.28, 0.14)	-0.06 (-0.24, 0.11)	0.05 (-0.10, 0.20)	0.06 (-0.07, 0.18)	referent	
P-value compared to referent	0.53	0.48	0.53	0.37	referent	
Global Composite Score						
Adjusted Z-score, mean (SE)	0.01 (0.04)	0.01 (0.03)	0.07 (0.03)	0.10 (0.03)	0.06 (0.04)	0.30
Difference from referent group, mean (95% CI)	-0.05 (-0.16, 0.07)	-0.04 (-0.14, 0.05)	0.01 (-0.07, 0.10)	0.04 (-0.03, 0.11)	referent	
P-value compared to referent	0.44	0.37	0.77	0.22	referent	

Positive changes in Z-score indicate improvement over time

Adjustments were based on repeated measure mixed-effect model assessing the least square means and comparing each LDL-C category with LDL-C ≥ 2.6 mmol/L. Model is adjusted for age, sex, education level, treatment, randomization strata, baseline LDL-C values, baseline statin intensity, baseline Z-scores, visit, and treatment by visit interaction term.

Table S5. Patient Self-Report of Change in Memory and Executive Function Since the Start of the Trial

Everyday Cognition Domain	LDL-C at Four Weeks in mmol/L					P _{trend}
	<0·5 (N=2669)	0·5-1·3 (N=8003)	1·3-1·8 (N=3444)	1·8-2·6 (N=7471)	≥ 2·6 (N=4395)	
Memory						
No. of patients	2304	6742	2771	6122	3541	0·11
Mean (SD)	1·17 (0·38)	1·15 (0·36)	1·18 (0·41)	1·16 (0·37)	1·18 (0·40)	
Adjusted P-value vs referent	0·61	0·0452	0·18	0·08	referent	
Executive Functioning, Total						
No. of patients	2304	6741	2769	6120	3541	0·12
Mean (SD)	1·12 (0·32)	1·10 (0·29)	1·13 (0·34)	1·11 (0·31)	1·13 (0·34)	
Adjusted P-value vs referent	0·81	0·15	0·53	0·21	referent	
Subdomains of Executive Function						
Planning						0·27
No. of patients	2302	6737	2767	6119	3539	
Mean (SD)	1·11 (0·34)	1·08 (0·28)	1·12 (0·35)	1·09 (0·31)	1·12 (0·35)	
Adjusted P-value vs referent	0·53	0·0373	0·97	0·73	referent	
Organization						0·98
No. of patients	2302	6741	2768	6113	3538	
Mean (SD)	1·10 (0·31)	1·09 (0·30)	1·12 (0·34)	1·09 (0·32)	1·12 (0·35)	
Adjusted P-value vs referent	>0·99	0·69	0·50	0·70	referent	
Divided Attention						0·0374
No. of patients	2302	6726	2763	6106	3533	
Mean (SD)	1·15 (0·40)	1·13 (0·36)	1·17 (0·42)	1·14 (0·38)	1·17 (0·42)	
Adjusted P-value vs referent	0·39	0·0472	0·96	0·22	referent	
Total Score						0·0168
No. of patients	2304	6743	2771	6122	3541	
Mean (SD)	1·13 (0·32)	1·12 (0·29)	1·15 (0·35)	1·13 (0·31)	1·15 (0·35)	
Adjusted P-value vs referent	0·49	0·019	0·94	0·0319	referent	

Table S6. Primary and Key Secondary Endpoint and Adverse Events in Patients with Ultra-Low LDL-C at Four Weeks

	LDL cholesterol in mmol/L (mg/dL) at four weeks						
	<0.26 (<10 mg/dL, N=504)			<0.4 (<15 mg/dL, N=1335)			≥2.6 (≥ 100 mg/dL, N=4395)
	n (%)	Adjusted HR (95% CI)	P*	n (%)	Adjusted HR (95% CI)	P*	n (%)
Efficacy Endpoints							
Primary efficacy endpoint	37 (7.3)	0.69 (0.49-0.97)	0.0354	105 (7.9)	0.71 (0.56-0.89)	0.0031	521 (11.9)
CV death, MI, stroke	22 (4.4)	0.59 (0.37-0.92)	0.0203	66 (4.9)	0.66 (0.50-0.88)	0.0049	345 (7.8)
	n (%)	Adjusted OR (95% CI)	P*	n (%)	Adjusted OR (95% CI)	P*	n (%)
Safety Endpoints							
Serious adverse events	115 (22.8)	0.94 (0.74-1.20)	0.61	313 (23.4)	0.96 (0.81-1.13)	0.63	1022 (23.3)
Adverse events leading to drug discontinuation†	18 (3.6)	1.08 (0.63-1.85)	0.78	42 (3.1)	0.89 (0.60-1.32)	0.56	149 (3.4)

*P value compared to the group achieving an LDL-C ≥ 2.6 mmol/L (≥ 100 mg/dL) at four weeks.

† Excludes injection site reactions

There were too few of the other eight prespecified safety events for an adjusted analysis in the two ultra-low LDL-C groups.